

**Md Hafiz All Hosen¹, Koushik Sarker¹,
Mousumi Sarker Chhanda^{2*} and Nipa
Gupta²**

¹Department of Fisheries Management, Faculty of Fisheries, Hajee Mohammad Danesh Science and Technology University, Bangladesh

²Department of Aquaculture, Faculty of Fisheries, Hajee Mohammad Danesh Science and Technology University, Bangladesh

Received: 16 May, 2019

Accepted: 02 August, 2019

Published: 05 August, 2019

***Corresponding author:** Mousumi Sarker Chhanda, Assistant Professor, Department of Aquaculture, Hajee Mohammad Danesh Science and Technology University (HSTU), Dinajpur-5200, Bangladesh, Tel: 01719471799; Email: chh.sarker@hstu.ac.bd

Keywords: Depth; Growth performance; Indian major Carps; Poly culture system

<https://www.peertechz.com>



Research Article

Effects of water depth on growth performance of Indian major carps at a poly culture system in Bangladesh

Abstract

The main aim of this study was to assess the effect of water depth on the growth performance (length and weight) of table size Indian major carp (*Gibelion catla*, *Labeo rohita*, and *Cirrhinus mrigala*) fishes. Fishes were recorded under different depth (1.20 m, 1.80 m and 2.80 m) for about eight months from February 2016 to September 2016. Water depth was considered as treatment 1 (T_1) for 1.20 m, treatment 2 (T_2) for 1.80 m and treatment 3 (T_3) for 2.80 m, respectively. Randomized Complete Block Design (RCBD) with two replications of each treatment was differentiated at significant level 0.05. Various water quality parameters such as water temperature, dissolve oxygen, transparency, and pH were also recorded. The survival rate was found maximum ($99.23 \pm 0.4\%$) for *Labeo rohita* fish at 2.80 m water depth and lowest ($95 \pm 2.1\%$) survival rate was observed for *Gibelion catla* fish at 1.20 m water depth. Results of this study showed that *Gibelion catla*, *Labeo rohita* and *Cirrhinus mrigala* fish cultivated at four feet depth of water produces lowest average weight of 1398 g, 1048 g and 1050 g, respectively with an average length of 47 cm, 49.2 cm and 46.4 cm, respectively. The maximum average weight of fish 3667 g, 2700 g and 2800 g were obtained at 2.80 m depth of water with the maximum average length of 63.2 cm, 60.0 cm and 58.6 cm for *Gibelion catla*, *Labeo rohita* and *Cirrhinus mrigala* fish, respectively. From this study, it can be observed that the fishes cultivated at low depth of water yielded lowest, and shorter in length in contrast to the carps cultivated at higher depth of pond water. However, growth of *Gibelion catla* was notably high among other carps.

Introduction

Bangladesh is one of the resourceful countries in the South-East Asia where different types of fish species are cultivated for fulfilling high quality protein requirement. Carp fish is one of the most significant fish species for aquaculture all over the world and represent the species of choice due to its high growth rate, ease in reproduction, tolerance to environmental stress and its market demand. Though there are at least 265 freshwater fish species in the country [1]. But only 4 native and 12 exotic carp species are cultured in Bangladesh [2]. Usually, large carp including Rui (*Labeo rohita*), Mrigel (*Cirrhinus mrigala*), Catla (*Gibelion catla*), grass carp and silver carp are common aquaculture species in Bangladesh [3,4].

Fish naturally tend to select the habitat that is most suitable for their physiological requirements. This behavior is known as 'habitat selection' or 'enviro regulation' [5]. The optimum fish production is totally dependent on the physical, chemical and biological qualities of water to most of the extent. Hence, successful pond management requires an understanding of water quality. Water quality is determined by variables like

temperature, transparency, turbidity, water color, carbon dioxide, pH, alkalinity, hardness, unionised ammonia, nitrite, nitrate, primary productivity, BOD and plankton population [6]. In addition to water quality ponds depth also significantly affect the growth rate and survival on fish cultivation [5].

Several researches have already carried out some researches regarding effect of water quality and depth of ponds on fish growth rate. The optimum pH range differs among species; however, the pH 6.5–9.0 range is generally accepted for fish culture [7]. The temperature at which the best growths of most carp for the tropical is 28–32°C [8]. According to Banerjee (1967) dissolve oxygen (DO) for the good production of fish should be above 5ppm [9]. Clerk (1986) reported that BOD range of 2 to 4 mg L⁻¹ does not show pollution while levels beyond 5 mg L⁻¹ are indicative of serious pollution [10]. According to Boyd and Lichtkoppler (1979), fishes are avoided free CO₂ levels as low as 5 mg L⁻¹, but most species can survive in waters containing up to 60 mgL⁻¹ carbon dioxide, provided DO concentrations are high [11]. Ali et al., (2013) pointed that the effect of water depth on Nile tilapia fingerlings and adults significantly affect the growth performance and survival rate [5].

However, the effect of pond depth on the growth of this three valuable carp fish has not yet been done so far. Therefore, this study was conducted to evaluate the effect of pond depth on the growth performance of Indian major carp's fishes (*Gibelion catla*, *Labeo rohita* and *Cirrhinus mrigala*).

Materials and Methods

Experimental site, experimental design and pond facilities

The experiment was carried out for a period of eight months during 1st February to 30th September, 2016 in six different ponds located at Niamatpur, Thakurgaon (Figure 1). The experiment had 3 treatments with 2 replications for each. Water depth was considering as treatment (T_1) for 1.20 m, treatment (T_2) for 1.80 m, treatment (T_3) for 2.8 m following randomized complete block design (RCBD). The ponds were rectangular in shape and the total surface area of these ponds was 650 decimals. Each pond had an inlet for watering but no outlet, free from aquatic vegetation, well-exposed to sunlight and sandy loam bottom. The stocked fish fed at morning and afternoon daily at the rate of 5% body weight with commercial pellet feed (30% crude protein) of name Nourish Poultry and Hatchery Ltd.

Pond preparation

Before starting the experiment, the ponds were dried and made free from aquatic vegetation. After drying, liming (CaO) was done in all the ponds at the rate of 250 kg/hectare. Ponds were then filled with ground water at required depth. Seven days after liming, Urea and triple super phosphate (TSP) were applied each at the rate of 39.0 and 37.0 kg/hectare, respectively. Seven days after fertilization, sumithion was applied in all the ponds at the rate 25 Liter/hectare.

Stocking

Experimental fishes were stocked in all ponds in the morning on 2nd February following the experimental design. Containers were kept floating in water about 30 min in the experimental ponds for acclimatization of temperature before releasing the carp fishes. The weight of approximately 10% of each species for each pond was measured and recorded for estimating initial stocking biomass as well as adjust feeding rate of fishes. Table 1 shows the stocking density in different ponds.

Post stocking management

Fertilization: Fertilization was done with Urea (50 kg/hectare) and TSP (25 kg/hectare) at fortnightly basis during the study period. TSP was dissolved in water for 24 hours before and Urea was dissolved in a bucket in the morning and then applied by spreading with a mug on the pond surface. Fertilization was done fortnightly.

Study of physicochemical parameters of water: The physico-chemical parameters of pond water were recorded fortnightly throughout the experimental period between 10 A.M. and 12 A.M. Physical parameters such as; water temperature (°C), transparency (cm), and water depth (m), dissolved




Figure 1: Experimental Location on the map of ThakurgaonSadarUpazila of Thakurgaon District.

Table 1: Stocking density were assigned in different ponds (nos = number of fishes).

Experimental fish	Stocking density (fish/decimal)
<i>Gibelion catla</i>	2 nos.
<i>Labeo rohita</i>	5 nos.
<i>Cirrhinus mrigala</i>	5 nos

oxygen (mg/L) and pH were measured at the pond site on each sampling day. Depth of water of the experimental ponds was estimated with the help of a graduated wooden scale.

Growth parameters:

Weight gain (g)

The weight gain was calculated through the following equation:

$$\text{Weight gain (g)} = \text{Mean final weight (g)} - \text{Mean initial weight (g)} \quad (1)$$

Specific growth rate

Specific growth rate of fish under different treatments was calculated using the formula:

$$\text{SGR (\%)} = \frac{\ln W_2 - \ln W_1}{T} \times 100 \quad (2)$$

where, $\ln W_2 - \ln W_1$ is the difference of logarithm of initial and final weight and T is the duration of the experiment (days).

Survival rate

The survival rate (SR) of fish was calculated as follow

$$\text{SR (\%)} = \frac{\text{Final total number of fish}}{\text{Initial total number of fish}} \times 100 \quad (3)$$

Statistical analysis

The data were analyzed by Statistical Analysis System (SPSS Windows version 22). Treatment means were compared at $p < 0.05$ according to the Duncan's New Multiple Range Test (DNMRT) and Least Significant Difference (LSD).

Approval

This experiment was performed under the Department of Fisheries Management in accordance with university rules and

regulations. Approved and revised by the post graduate defence committee of Hajee Mohammad Danesh Science & Technology University, Dinajpur-5200, Bangladesh (approval and certified no. 914).

Results and Discussion

1. Analysis of physicochemical Parameters

Temperature

The monthly values of water temperature throughout the experimental period are shown in table 2. It was found that, between the temperature range of 18–30°C, the growth of the fishes was high. This finding was concurred with Nazish and Mateen (2010) as they reported that the freshwater fish have an optimum growing temperature in the range of 25–30°C at which they grow quickly [12]. However, there was no significant difference in temperature of water in the month of April and May. In July lowest temperature was recorded at 18°C while the highest temperature (30°C) was noted in May. Similar findings in temperature of pond water were reported by Nwipieet al. (2015), Kohinoor *et al.* (2012) [13, 14].

Dissolved oxygen (DO)

The change of dissolved oxygen in pond at different depth of water is shown in table 3. Low dissolve oxygen (5.4 ppm) was found in the month of February at T₁ treatment while maximum dissolve Oxygen (9.07 ppm) was found in the month of July at T₃ treatment. Similar findings were also reported by Bhatnagar and Singh (2010) that DO level greater than 5 ppm is essential to support good fish production [8].

Transparency

The transparency values throughout the experiment were shown in table 4. Low transparency (28.12 cm) was recorded at T₃ treatment in February while high transparency (34.3 cm) was at T₂ treatment in July. This result was in good agreement with Boyd (1982) that the transparency between 25 to 40 cm as appropriate for fish culture [15]. However, Azim and Wahab (2003) reported the transparency value as 36.2 cm in weed based carp polyculture pond [16].

pH

The water pH was measured during the whole experiment was shown in table 5. Analysis of the data revealed that within the same treatment the change of pH was not significant from February to May for all the treatment. The highest pH value (8.6) was recorded at T₃ treatment in the month of February while the lowest pH value (6.7) was recorded at same treatment in the month of July. From the experiment, it was observed that the maximum growth of fish occurred at a pH range of 6.7 to 8.6. The similar observations have been made by Singh [17].

Growth parameters

Survival rate (SR) of carp fishes: The survival rate of *Gibelion catla*, *Labeo rohita* and *Cirrhinus mrigala* fishes are presented in figures 2–4, respectively. It was found that the effect of water

Table 2: Monthly Temperature Variation (mean±SD) of Pond Water under Different Treatments.

Month	Treatments		
	T ₁	T ₂	T ₃
February	22.30±0.2 ^c	21.50±0.5 ^c	20.00±1 ^c
March	25.93±0.06 ^b	24.63±0.6 ^{ab}	23.33±1.53 ^b
April	29.30±0.25 ^a	28.10±0.40 ^a	26.00±1.0 ^a
May	29.90±0.10 ^a	28.67±.30 ^a	27.33±.57 ^a
June	23.53±.50 ^b	22.53±0.42 ^c	20.67±.60 ^b
July	20.10±0.1 ^c	18.67±0.3 ^c	18.00±1.0 ^b
August	22.50±0.1 ^c	21.10±.36 ^c	19.00±1.0 ^c
September	27.77±0.20 ^{ab}	26.13±0.35 ^b	24.50±0.5 ^c

^{a-c}mean followed by different subscript alphabet in each raw significantly different (p<0.05) among different temperature within same treatment.

Table 3: Monthly Dissolved Oxygen Variations (mean±SD) of Pond Water under Different Treatments.

Month	Treatments		
	T ₁	T ₂	T ₃
February	5.50±0.1 ^c	5.80±0.1 ^b	6.10±0.10 ^b
March	6.20±1.52 ^b	6.50±0.05 ^b	6.67±0.57 ^b
April	6.33±0.15 ^b	6.87±0.006 ^b	7.20±0.05 ^a
May	6.20±0.0.10 ^b	6.53±0.57 ^b	7.06±0.06 ^a
June	6.50±0.10 ^b	6.90±0.0.10 ^b	7.33±0.57 ^a
July	7.03±0.57 ^a	7.70±0.10 ^a	8.50±0.57 ^a
August	6.90±0.10 ^a	7.60±0.10 ^a	8.33±0.06 ^a
September	6.20±0.10 ^c	6.40±0.20 ^b	6.76±0.057 ^b

^{a-c}mean followed by different subscript alphabet in each raw significantly different (p<0.05) among different dissolved oxygen values within same treatment.

Table 4: Monthly Transparency Variations (mean ± SD) of Pond Water under Different Treatments.

Month	Treatments		
	T ₁	T ₂	T ₃
February	28.13±0.15 ^c	28.16±0.05 ^c	27.97±0.15 ^c
March	29.30±0.10 ^b	29.33±0.06 ^b	29.16±0.15 ^b
April	29.10±0.1 ^b	29.40±0.1 ^b	29.20±0.06 ^b
May	28.20±0.10 ^c	28.06±0.11 ^c	28.10±0.1 ^b
June	29.96±0.15 ^b	30.40±0.10 ^b	30.20±0.05 ^a
July	34.06±0.12 ^a	34.20±0.10 ^a	34.23±0.05 ^a
August	32.50±0.10 ^a	32.27±0.05 ^a	32.76±0.49 ^a
September	29.20±0.1 ^b	29.96±0.15 ^b	29.10±0.10 ^b

^{a-c}mean followed by different subscript alphabet in each raw significantly different (p<0.05) among different transparency values within same treatment.

Table 5: Monthly pH Variations (mean ± SD) of Pond Water under Different Treatments.

Month	Treatments		
	T ₁	T ₂	T ₃
February	8.43±0.11 ^a	8.50±0.1 ^a	8.50±0.10 ^a
March	8.23±0.06 ^a	8.23±0.05 ^a	8.10±0.10 ^a
April	8.40±0.1 ^a	8.30±0.10 ^a	8.20±0.10 ^a
May	8.06±0.06 ^a	7.90±8.1 ^a	8.10±0.10 ^a
June	8.17±0.15 ^a	7.90±0.10 ^a	7.70±0.10 ^{ab}
July	7.60±0.10 ^b	7.10±0.10 ^b	6.60±0.10 ^c
August	7.80±0.10 ^{ab}	7.50±0.10 ^b	6.67±0.55 ^c
September	8.10±0.10 ^a	8.00±0.10 ^a	8.00±0.12 ^a

^{a-c}mean followed by different subscript alphabet in each raw significantly different (p<0.05) among different pH values within same treatment.

depth on survival rate was remained significant ($p < 0.05$). At the end of experiment, the highest survival rate (99.23%) was observed in T_3 followed by T_2 (99.0%) and the lowest survival rate (95.0%) was recorded in T_1 for *Gibelion catla* fishes in figure 2. It was also observed that the survival rate was maximum at T_3 and lowest at T_1 as depicted in figure 2. However, maximum survival (99.23%) rate was found for *Labeo rohita* fishes at T_3 as shown in figure 3. The survival rate of this present study was corresponding to Miah et al., [18] and found the survival rate from 89.23-99.23% and similar findings were also reported by Mamun and Mahmud [19].

Specific growth rate (SGR) of carp fishes: Specific growth rate of *Gibelion catla*, *Labeo rohita* and *Cirrhinus mrigala* fishes are presented in figures 5-7 respectively. It was found that the effect of water depth on specific growth rate was remained significant ($p < 0.05$). The specific growth rate of *Gibelion catla*, *Labeo rohita* and *Cirrhinus mrigala* fishes in this experiment reveals that at the early stage of rearing the growth of fishes were high. The specific growth rate decreases at the end of the experiment for all the treatment (T_1 , T_2 and T_3). However, specific growth rate at T_3 was higher while compare to the T_1




Figure 2: Survival rate of *Gibelion catla* fish at different water level.




Figure 3: Survival rate of *Labeo rohita* fish at different water level.




Figure 4: Survival rate of *Cirrhinus mrigala* fish at different water level.




Figure 5: Monthly variations of specific growth rate of *Gibelion catla* fish.




Figure 6: Monthly variations of specific growth rate of *Labeo rohita* fish.




Figure 7: Monthly variations of specific growth rate of *Cirrhinus mrigala* fish.

and T_2 for all carp fishes. Similar findings were also reported by Mamun and Mahmud [19].

Average weight and length gain of carp fishes: Average weight and length gain of *Gibelion catla*, *Labeo rohita* and *Cirrhinus mrigala* fishes are presented in table 6. The individual average weight (2511 ± 95.4 gm) gain was maximum for *Gibelion catla* fishes at T_3 and minimum individual average weight (769 ± 43 gm) gain was found for *Cirrhinus mrigala* fishes at T_1 treatment. Analysis of the data also reveals that the individual average length gain was maximum (53.9 ± 9.3 cm) for *Gibelion catla* fishes at T_3 treatment and minimum (39 ± 6.2 cm) individual average length gain was found for *Labeo rohita* fish at T_1 treatment (Table 5). The average specific growth was found maximum ($1.12 \pm 0.1\%$) for *Labeo rohita* fishes at T_3 treatment while lowest ($0.62 \pm 0.02\%$) specific growth rate was found for *Cirrhinus mrigala* fish at T_1 treatment. Analyzing of the data also reveals that survival rate of all table sizes carp fishes was found greater than 95%. However, the survival rate was found

Table 6: Growth performance of *Gibelion catla*, *Labeo rohita* and *Cirrhinus mrigala* fish (Mean±SD) in different treatments over the experimental periods.

Species		Item	Treatments		
			T ₁	T ₂	T ₃
<i>Gibelioncatla</i>	Initially	No. of fish (fish/hectare)	494±0	494±0	494±0
		Av. Individual weight of fish (gm/fish)	250±10	250±12	250±9
		Av. Individual total length of fish (cm/fish)	20±1.5	20±1.8	20±2.3
	Finally	No. of fish (fish/hectare)	469±0	484±0	485±0
		Av. Individual weight of fish (gm/fish)	979.50±80 ^c	1813.60±82.2 ^b	2511±1156 ^a
		Body weight gain (gm/fish)	729±78 ^c	1563.1±92 ^b	2260±1100 ^a
		Av. Individual total length of fish (cm/fish)	39.60±7.3 ^c	48.20±8 ^b	53.90±9.3 ^a
		Body length gain (cm/fish)	19.65±6.5 ^c	28.7±7 ^b	32±8.5 ^a
		Specific growth rate (%growth)	0.8±0.03 ^b	1.1±0.08 ^a	1.19±0.7 ^a
Survival rate (%)	95±2.1 ^{ab}	98.3±3.2 ^a	98±0.8 ^a		
<i>Labeorohita</i>	Initially	No. of fish (fish/hectare)	1235±0	1235±0	1235±0
		Av. Individual weight of fish (gm/fish)	250±7	250±11	250±8
		Av. Individual total length of fish (cm/fish)	21±0.6	21±0.5	22±0.7
	Finally	No. of fish (fish/hectare)	1211±0	1216±0	1223±0
		Av. Individual weight of fish (gm/fish)	759.7±288 ^c	1307±558 ^b	1862.08±838 ^a
		Body weight gain (gm/fish)	508±92 ^c	1050±81 ^b	1621±732 ^a
		Av. Individual total length of fish (cm/fish)	39±6.2 ^c	46±7.4 ^b	50.54±9 ^a
		Body length gain (cm/fish)	17±0.41 ^c	23.8±6.8 ^b	28.50±7.1 ^a
		Specific growth rate (%growth)	0.63±0.04 ^c	0.9±0.008 ^b	1.12±0.1 ^a
Survival rate (%)	98.2±0.3 ^{ab}	99.0±0.1 ^a	99.23±0.4 ^a		
<i>Cirrhinusmrigala</i>	Initially	No. of fish (fish/hectare)	1235±0	1235±0	1235±0
		Av. Individual weight of fish (gm/fish)	250±9	250±11	250±10
		Av. Individual total length of fish (cm/fish)	21.40±0.5	20.80±0.8	22±0.2
	Finally	No. of fish (fish/hectare)	1186±0	1208±0	1215±0
		Av. Individual weight of fish (gm/fish)	759.70±273 ^c	1307.70±528 ^b	1862±793 ^a
		Body weight gain (gm/fish)	500.2±92 ^c	1058±52 ^b	1610±725 ^a
		Av. Individual total length of fish (cm/fish)	41±5.4 ^c	45.80±5.3 ^b	51.30±7.3 ^a
		Body length gain (cm/fish)	19.50±3.8 ^c	24±4.1 ^b	30.10±6.2 ^a
		Specific growth rate (%growth)	0.62±0.02 ^b	0.85±0.04 ^a	1.10±0.07 ^a
Survival rate (%)	96.00±2.3 ^b	97.80±1.5 ^a	98.40±1.1 ^a		

^{a-c}mean followed by different subscript alphabet in each row significantly different (p<0.05) among different species between each treatment.

maximum (99.23±0.4%) for *Labeo rohita* fish at T₃ treatment and lowest (95±2.1%) survival rate was found for *Gibelion catla* fishes at T₁ treatment.

Effects of Water Depth on the Weight and Length of Carp Fish

Effect of water depth on the weight of *Gibelion catla* fish: Analysis of the data showed that the weight of *Gibelion catla* fish were significantly (p<0.05) affected by the single effect of water depth. It was observed that *Gibelion catla* fish cultivated on 2.80 m depth of water yielded higher fish weight while compare with the *Gibelion catla* fish cultivated at the lower water depth. Figure 8 showed that the mean weight of *Gibelion catla* fish increased linearly from 1.20 m to 1.80 m depth of water and from 1.80 m to 2.80 m depth of water mean weight of *Gibelion catla* fish decreased slightly as compared to 1.20 m to 1.8 m depth of water. Therefore, it was concluded that carp fish cultivated at higher water depth returned higher amount of fish weight in contrast to the fish cultivated at lower water depth. This result was in great concord with Ali et al. who




Figure 8: Weight of *Gibelion catla* fish at various water depths.

reported that the weight of Nile Tilapia is increased with the increase of water depth [5]. Similar findings were also reported by Stoll et al., [20].

Effect of water depth on the length of *Gibelion catla* fish:

The effect of water depth on the length of *Gibelion catla* fish found significant ($p < 0.05$). The length of the *Gibelion catla* fish was found maximum when it was cultivated at 2.80 m depth of water and minimum was found while cultivation of fish was done at 1.20 m depth of water. Figure 9 exhibits that the length of *Gibelion catla* fish increased linearly from 1.20 m depth of water to 1.80 m depth of water but slightly decreased from 1.80 m to 2.80 m depth of water. Therefore, it was concluded that carp fish cultivated at higher depth (2.80 m) of water generates fish of larger in length while compare with carp fish cultivated at lower depth (1.20 m) of water yielded fish of smaller in length. The result was in agreement with Ali *et al.* (2013) who reported that the length of Nile Tilapia is increased with the increase of water depth [5]. Similar observations were also Takashi and Tadashi [21].

Effect of water depth on the weight of *Labeo rohita* fish: Significant effect of water depth was observed in respect of weight (g) of *Labeo rohita* fish yield ($p < 0.05$). It was observed that the weight of *Labeo rohita* fish was gradually increased with the increase of depth of water level. The highest weight of *Labeo rohita* fish was recorded when fish was cultivated at 2.80 m depth of water while lowest weight of *Labeo rohita* fish was recorded at 1.20 m depth of water. Figure 10 showed that the mean weight of *Labeo rohita* fish increased linearly from 1.20 m to 2.80 m depth of water. Therefore, it was concluded that carp fish cultivated at higher water depth (2.80 m) returned higher amount of fish weight in contrast to the fish cultivated at lower water depth (1.20 m). Similar findings were also pointed by Payne, et al., [22].

Effect of water depth on the length of *Labeo rohita* fish: Analysis of the data revealed that the depth of water significantly ($p < 0.05$) affected the length of *Labeo rohita* fish. As the cultivation of fish was done at the low water depth, the length of the *Labeo rohita* fish yielded minimum value. Result showed that the *Labeo rohita* fish cultivated at 2.80 m depth of water yielded maximum mean length over the cultivation of *Labeo rohita* fish at 1.20 m. Figure 11 revealed that the length of *Gibelion catla* fish increased linearly from 1.20 m depth of water to 1.80 m depth of water but slightly decreased from 1.80 m to 2.80 m depth of water in contrast to 1.20 m to 1.80 m depth of water. Therefore, it was concluded that carp fish cultivated at




Figure 9: Length of *Gibelion catla* fish at various water depths.




Figure 10: Weight of *Labeo rohita* fish at various water depths.




Figure 11: Length of *Labeo rohita* fish at various water depths.

higher depth (2.80 m) of water produced fish of larger in length while compare with carp fish cultivated at lower depth (1.20 m) of water. This result was full coincided with the reported values of Stoll et al. that the depth of ponds should not be less than 100–200 cm for the maximum growth of tilapia fish [20].

Effect of water depth on the weight of *Cirrhinus mrigala* fish: Significant effect of water depth on the weight of *Cirrhinus mrigala* fish was observed ($p < 0.05$). It was observed that the weight of *Cirrhinus mrigala* fish increased gradually as the depth of water increases. The highest weight (g) of *Cirrhinus mrigala* fish was noted when it was cultivated on pond with a constant 2.80 m depth of water and the lowest fish weight was recorded in correspond to the 1.20 m depth of water. Figure 12 showed that the mean weight of *Cirrhinus mrigala* fish increased linearly from 1.20 m to 2.80 m depth of water. Therefore, it was recommended that carp (*Cirrhinus mrigala*) fish cultivated at higher water depth (2.80 m) returned higher amount of fish weight in contrast to the carp fish cultivated at lower water depth (1.20 m). This result was in great concord with Ali et al., (2013) who reported that the weight of Nile Tilapia is increased with the increase of water depth [5]. Similar conclusion was also reported by Takashi and Tadashi and Stoll et al., [20,21].

Effect of water depth on the length of *Cirrhinus mrigala* fish: The effect of water depth on the length of *Cirrhinus mrigala* fish found significant ($p < 0.05$). The length of the *Cirrhinus mrigala*

fish was found maximum when it was cultivated at 2.80 m depth of water and minimum was found while cultivation of paddy was done at 1.20 m depth of water. Figure 13 showed that the mean weight of *Cirrhinus mrigala* fish increased linearly 1.20 m to 2.80 m depth of water. Therefore, it was concluded that carp fish cultivated at higher depth (2.80 m) of water generates fish of larger in length while compare with carp fish cultivated at lower depth (1.20 m) of water yielded fish of smaller in length. The result was in full agreement with Ali et al., (2013) who reported that the length of Nile Tilapia is increased with the increase of water depth [5]. Similar outcomes were also noted by Takashi and Tadashi and Stoll et al., [20,21].




Figure 12: Weight of *Cirrhinus mrigala* fish at various water depths.




Figure 13: Length of *Cirrhinus mrigala* fish at various water depths.

Conclusions

Therefore, the outcome of this study suggests that the carp fish cultivated at 2.80 m depth of water will be beneficial and more economical for the farmer and more economy will be earned in compared to the 1.20 m and 2.80 m depth of water.

Author Contributions

Md. Hafiz All Hosen was the principal investigator of this experiment and was analyzed and writing the original manuscripts, Koushik Sarker conducted the experiment and

analyzes the data, Mousumi Sarker Chhanda and Nipa Gupta investigated the experiment and wrote the manuscripts.

References

- Rahman AKA (2005) Freshwater Fishes of Bangladesh. Second edition, Zoological Society of Bangladesh, Dhaka, Bangladesh 394. [Link: http://bit.ly/319jsy8](http://bit.ly/319jsy8)
- (2012) National Fish Week 2012 Compendium. Department of Fisheries, Ministry of Fisheries and Livestock, Dhaka, Bangladesh 144.
- Hossain MY, Rahman MM, Jasmin S, Ibrahim AHM, Ahmed ZZ, et al. (2008) Comparison studies on water quality and plankton production between perennial and non-perennial ponds in Bangladesh. Journal of Fisheries and Aquatic Science 3: 176-183. [Link: http://bit.ly/2YFCGtM](http://bit.ly/2YFCGtM)
- Rahman MM, Verdegem MCJ, Wahab MA (2008) Effects of tilapia (*Oreochromis nilotica* L.) addition and artificial feeding on water quality, and fish growth & production in Labeorohita-common carp bi-culture ponds. Aquaculture Research 39: 1579-1587. [Link: http://bit.ly/2Zwepro](http://bit.ly/2Zwepro)
- Ali MAM, El-Feky AMI, Khouraiha HM, El-Sherif MS (2013) Effect of water depth on growth performance and survival rate of mixed sex Nile tilapia fingerlings and adults. The Egyptian Society of Animal Production, 50: 194-199. [Link: http://bit.ly/2Yldset](http://bit.ly/2Yldset)
- Anita Bhatnagar, Pooja Devi (2013) Water quality guidelines for the management of pond fish culture. International journal of environmental sciences 3: 1981-2009. [Link: http://bit.ly/2YF3Tg3](http://bit.ly/2YF3Tg3)
- Zweig RD, Morton JD, Stewart MM (1999) Source water quality for aquaculture: A guide for assessment The World Bank, Washington 62. [Link: http://bit.ly/2Kg0RuZ](http://bit.ly/2Kg0RuZ)
- Bhatnagar A, Singh G, (2010) Culture fisheries in village ponds: a multi-location study in Haryana, India. Agriculture and Biology Journal of North America, 1: 961-968. [Link: http://bit.ly/2ZuUIQu](http://bit.ly/2ZuUIQu)
- Banerjea SM (1967) Water quality and soil condition of fishponds in some states of India in relation to fish production, Indian journal of fisheries. Open Journal of Applied Sciences 14: 115-144. [Link: http://bit.ly/2ZuoncN](http://bit.ly/2ZuoncN)
- Clerk RB (1986) Marine Pollution. Fifth Edition, Oxford, 256.
- Boyd CE, Lichtkoppler F (1979) Water Quality Management in Fish Ponds. Research and Development Series No. 22, International Centre for Aquaculture (J.C.A.A) Experimental Station Auburn University, Alabama, 45-47.
- Nazish N, Mateen A (2010) Winter Growth of Carps under Different Semi-Intensive Culture Conditions. Pakistan Veterinary Journal, 31: 134-136. [Link: http://bit.ly/334FBzv](http://bit.ly/334FBzv)
- Nwipie GN, Erondu ES, Zabbey N, (2015) Influence of Stocking Density on Growth and Survival of Post Fry of the African Mud Catfish, *Clarias gariepinus*. Fish Aquac J 6: 1. [Link: http://bit.ly/338KAPG](http://bit.ly/338KAPG)
- Kohinoor AHM, Khan MM, Yeasmine S, Mandol P, Islam MS (2012) Effects of stocking density on growth and production performance of indigenous stinging catfish, *Heteropneustes fossilis* (Bloch) Int J Agril Res Innov Tech 2: 9-14. [Link: http://bit.ly/33byjKC](http://bit.ly/33byjKC)
- Boyd CE (1982) Water quality management of pond fish culture. Elsevier Sci Pub Co Amsterdam-Oxford, New York, 318. [Link: http://bit.ly/2MKQrdh](http://bit.ly/2MKQrdh)
- Azim ME, Wahab MA (2003) Development of a duckweed-fed carp polyculture system in Bangladesh. Aquaculture, 18: 425-438. [Link: http://bit.ly/2GLQUDD](http://bit.ly/2GLQUDD)
- Singh RK (2007) Fluctuation of plankton population and some hydrological factors of the Mc Pherson Lake, Allahabad.

18. Miah (1997) Effect of stocking ratios on the growth and production of fishes in mixed polyculture system. Bangladesh J Fish 20:135-138. [Link: http://bit.ly/2ZpNc9E](http://bit.ly/2ZpNc9E)
19. Mamun AA, Mahmud AS, (2014) Study on the growth performance and production of juvenile Indian major carps (*GibelioncatlaGibelioncatla*, *Labeorohita* and *Cirrhinuscirrhose*) and their hybrids. Journal of Fisheries and aquatic science 9: 92-108. [Link: http://bit.ly/2Yn9Yme](http://bit.ly/2Yn9Yme)
20. Stoll S, Fischer P, Klahold P, Scheifhacken N, Hofmann H, et al. (2008) Effects of water depth and hydrodynamics on the growth and distribution of juvenile cyprinids in the littoral zone of a large pre-alpine lake. Journal of Fish Biology,

72: 1001-1022. [Link: http://bit.ly/2KkXIIT](http://bit.ly/2KkXIIT)

21. Takashi K, Tadashi K (2003) Water Depth as a Factor Influencing Growth Rate and Size of the Japanese Endemic Crayfish *Cambaroides japonicus* (Crustacea: Decapoda). Japanese Journal of Benthology, 58: 11-15. [Link: http://bit.ly/2Mxz8XY](http://bit.ly/2Mxz8XY)
22. Payne AL, Ridgway J, Hamer JL (1988) The influence of salt (NaCl) concentration and temperature on the growth of *Oreochromis spilurus*, *O. mossambicus* and red tilapia hybrids. In: *The Second International Symposium on Tilapia in Aquaculture* (ed. by R.S.V. Pullin, T Bhukaswan, K, Tonguthai & J.L. Maclean), Department of Fisheries, Bangkok, Thailand and ICLARM, Manila, Philippines, 481-487.

Discover a bigger Impact and Visibility of your article publication with Peertechz Publications

Highlights

- ❖ Signatory publisher of ORCID
- ❖ Signatory Publisher of DORA (San Francisco Declaration on Research Assessment)
- ❖ Articles archived in worlds' renowned service providers such as Portico, CNKI, AGRIS, TDNet, Base (Bielefeld University Library), CrossRef, Scilit, J-Gate etc.
- ❖ Journals indexed in ICMJE, SHERPA/ROMEO, Google Scholar etc.
- ❖ OAI-PMH (Open Archives Initiative Protocol for Metadata Harvesting)
- ❖ Dedicated Editorial Board for every journal
- ❖ Accurate and rapid peer-review process
- ❖ Increased citations of published articles through promotions
- ❖ Reduced timeline for article publication

Submit your articles and experience a new surge in publication services
(<https://www.peertechz.com/submission>).

Peertechz journals wishes everlasting success in your every endeavours.

Copyright: © 2019 Hosen HA, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Citation: Md Hosen HA, Sarker K, Chhanda MS, Gupta N (2019) Effects of water depth on growth performance of Indian major carps at a poly culture system in Bangladesh. Int J Aquac Fish Sci 5(3): 014-021. DOI: <http://doi.org/10.17352/2455-8400.000046>