

Research Article

Agronomic and yield performances of cucumber [*Cucumis sativus*] on soil amended with different rates of cattle manure in derived savannah Agroecological zone of Ondo state, Nigeria

Eleduma Ajayi Festus*

Department of Agricultural Technology, Rufus Giwa Polytechnic, PMB 1019, Owo, Ondo State, Nigeria

Received: 11 July, 2023

Accepted: 18 July, 2023


Published: 19 July, 2023

*Corresponding authors: Eleduma Ajayi Festus, Department of Agricultural Technology, Rufus Giwa Polytechnic, PMB 1019, Owo, Ondo State, Nigeria, Tel: +2348030416850; E-mail: eledumafestus@gmail.com

Keywords: Amend; Cattle manure; Cucumber; Inorganic fertilizer; Substitute

Copyright License: © 2023 Eleduma, A. F. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

<https://www.peertechzpublications.org>



Abstract

Most soils have experience nutrient depletion as a result of long-term farming, making good yields only possible with the application of fertilizers. A Randomized Complete Block Design (RCBD) was used to create a total of fifteen (15) plots in the study, which was conducted at the Rufus Giwa Polytechnic's Teaching Research and Commercial Farms in Owo, Ondo State, Nigeria. The study examined the impact of various rates of cattle manure on cucumber. The results revealed that the performance of cucumbers was significantly impacted by an increase in the rate of cattle manure, with 15 tha⁻¹ of cattle manure producing the best results in terms of growth and yield (3.45 cm, 73.46 cm, 171.22 cm², 16.83, 9.83 cm, 21.89 cm, 17.19 cm, and 5.14 kg as stem girth, vine length, leaf area, number of leaves, number of fruits, fruit weight, and fruit circumference) and the lowest from control. According to the study, applying cattle manure in the study area at a rate of 15 tha⁻¹ considerably improves cucumber growth and yield. In order to improve the physical and chemical properties of the soil and eliminate the need to apply inorganic fertilizer, farmers in the research area are advised to amend their soils with 15 tha⁻¹ cattle dung.

Introduction

One of the most significant vegetables in the Cucubitaceae family, cucumber (*Cucumis sativus* L.) is grown for its fruit, which is a rich source of vitamins and minerals. According to Eifediyi and Remison [1], it comes in second place to tomatoes in Western Europe and ranks fourth in Asia behind tomatoes, cabbage, and onion. However, the crop hasn't been ranked because of its sparse use in tropical Africa.

Due to its excellent nutritional and economic potential, cucumber cultivation is quickly gaining popularity throughout Nigeria [2]. This crop has numerous health advantages when consumed fresh, including important antioxidant, anti-inflammatory, and anti-cancer advantages [3].

Cucumbers' fiber-rich skin, high potassium and magnesium content, and ability to assist the body excrete uric acid all serve to lower blood pressure and support nutritional processes. This is especially helpful for people with arthritis. Cucumbers' magnesium content also helps to improve blood circulation while calming nerves and muscles.

The basic soil minerals were mined via continuous farming of tropical soils without the use of soil additives; these nutrients must be replaced to stop the loss of nutrients and preserve production. Due to low soil fertility, this was typically to blame for the low yields of cucumbers observed in the tropics. Infertile soils result in bitter and deformed fruits, which are frequently rejected by consumers, in addition to low production [4].

According to Wang, et al. [5], organic manure has

traditionally been used as a soil supplement to improve soil productivity, raise organic carbon content, microbes, crumb structure, the soil's nutritional status, and increase crop production. In addition to calcium, potassium, phosphorus, magnesium, salt, and magnesium, which are all vital for plant growth, cattle manure also contains a significant amount of nitrogen, making it a great organic fertilizer. Contrary to inorganic fertilizer, organic manure enriches the soil with organic matter, enhancing its structure, nutrient retention, aeration, capacity to hold moisture, and water infiltration [6]. Additionally, it contains vital nutrients that are linked to high photosynthetic activity, which encourages the growth of roots and vegetables [7].

In view of the quest for organically produced agricultural produce, this study was therefore designed to evaluate the growth and yield responses of cucumber to different rates of cattle manure under field conditions in derived savannah agro-ecological of Ondo State, Nigeria.

Materials and methods

Experimental site and land preparation

The study was carried out on a total area of 104 m² (13 m x 8 m) at the Rufus Giwa Polytechnic's Teaching, Research, and Commercial Farms in Owo, Ondo State, which is situated between latitudes 7° 12' N and 5° 12' E in Nigeria's Tropical Rain Forest Ecological Zone. An intense rainy season and a bimodal rainfall pattern are features of the experimental site. The used area had been continuously tilled for roughly three years with no evidence of the application of either inorganic or organic fertilizer. To make the soil level and smooth to form seed beds suitable for simple germination and crop establishment, the earth was mechanically plowed and harrowed using a tractor and then ground with a shovel.

Soil sample analysis

Core soil samples were collected using a soil auger at a depth of 0 cm – 15 cm from 10 different points on the experimental field before planting. The samples were bulked, air dried, and sieved through a 2 mm sieve prior to analysis for physicochemical properties. The composite samples were analyzed for pH using a 1:2 (soil: water) suspension [8]. Particle size distribution was determined using the hydrometer method [9]. Total nitrogen was evaluated by the Macro Kjeldahl method [10] and exchangeable cations (K, Ca, Mg, and Na) after extraction with NH₄OH (pH7) using a flame photometer [11]. K in the filtered extract was determined using an Atomic Absorption Spectrometer [12]. Available phosphorus was determined with the Bray I method (Bray and Kurtz, 1965), as modified by Olsen and Sommers [13]. Total organic carbon was determined by the Walkley-Black method (Nelson and Sommers, 1982). The outcome in Table 1 revealed that the experimental site was sandy loam soil textural categorization and acidic in nature. The soil fertility was critically low as exemplified by the low Organic matter (0.65 kg) basic nutrients and exchangeable cations.

Source of cattle manure and analysis

Manure from cattle was sourced from the teaching, research, and commercial farms at Rufus Giwa Polytechnic in Owo, Ondo State. Before use, the sample was subjected to typical routine laboratory analysis, crushed, and shade dried to assess the nutrient contents [14]. The analysis of the cattle dung revealed that it had an appreciable quantity of organic matter and basic nutrients, while the pH is slightly alkaline in nature (6.36) as presented in Table 2.

Experimental design and treatments

Five treatment levels were used in the experiment, and each level was replicated three times for a total of fifteen (15) plots. The experiment was set up using a Randomized Complete Block Design (RCBD). T₁ = 0tha⁻¹ of cattle dung (used as the control), T₂ = 5tha⁻¹ of cattle manure, T₃ = 10tha⁻¹ of cattle manure, T₄ = 15tha⁻¹ of cattle manure, and T₅ = 20tha⁻¹ of cattle manure were the treatment levels. The dimensions of each block were 12 x 2 m (24 m²), the plots were 2.2 m (four m²), and the discard between replicates and plots was 0.5 m. Two weeks before planting, the cattle dung was spread equally across the plots and worked into the soil with a hoe to promote further decomposition and nutrient release.

Table 1: Soil Physical and Chemical Properties Before Planting.

parameters	value
Particle size distribution	
Sand (%)	70
Silt (%)	6.8
Clay (%)	23.2
Textural class	Sandy loam
Soil pH (H ₂ O)	4.24
Organic Carbon (%)	0.38
Organic Matter (%)	0.65
Total Nitrogen (%)	0.14
Available P (ppm)	0.45
Exchangeable cations	
Na (comkg ⁻¹)	0.96
Ca (comkg ⁻¹)	1.70
K (comkg ⁻¹)	0.39
Mg (comkg ⁻¹)	6.80

Table 2: Nutrient composition of Cattle manure used.

Cattle Manure	Value
pH (H ₂ O)	6.36
O.C (%)	13.87
O.M (%)	23.90
N (%)	0.85
K (comkg ⁻¹)	4.62
Na (comkg ⁻¹)	3.91
Ca (comkg ⁻¹)	3.30
Mg (comkg ⁻¹)	8.90
Av. P (ppm)	1.41

Crop establishment and management

The National Institute for Horticultural Research and Training (NIHORT), located in Ibadan, Oyo State, Nigeria, provided the cucumber seeds for the experiment (Apulia semences variety). When the first weeding exercise was performed two (2) weeks after planting, the seeds were thinned to one seedling per stand, resulting in a population of 25 plants per plot and a total plant population of 375 plants on the entire experimental site. The seeds were sown at a rate of two seeds per hole with a spacing of 50cm by 50cm between and within rows. Regular weeding was done as and when it was necessary. To manage infestations of fruit flies and foliage insect pests, insecticide (Cypermethrin) was used at the prescribed dosage of 20 ml/15 liters per knapsack sprayer.

Data collection and statistical analysis

Five (5) stands per plot were randomly tagged for growth and yield parameter collection. Vine length (cm) was determined using measuring tape. The tape was used to measure the height of the plant from the level of the soil to the top of the shoot apex [15]. The number of leaves was determined by visually counting the number of leaves per plant (Edegbai and Oki, 2022). Stem girth was determined using measuring tape by putting the tape around the stem of the plant [15]. Leaf area (cm²) was determined by placing the leaf on graph paper. Yield attributes were determined by counting the number of cucumber fruits per plant, and the weight of harvested fruit (kg) was determined by weighing using a digital weighing balance. Fruit length (cm) was determined using a measuring tape to measure the length of the fruit from the base to the tip, while fruit circumference (cm) was determined by putting the measuring tape around the fruit. Data collected were subjected to two-way analysis of variance (ANOVA) procedures for Randomized Complete Block Design (RCBD) using the SAS statistical package Version 9 [16], and means were compared at 5% using the Duncan Multiple Range Test (DMRT).

Results

Effect of cattle manure on vegetative growth parameters of cucumber

According to the result in Table 3, the vegetative growth of cucumber is considerably ($p > 0.05$) influenced by the varied rates of cattle manure. The growth characteristics responded favorably to rising manure application rates. The development of stem girth ranged from 1.02 to 3.45 cm, with the thinnest girth (1.02 cm²) being measured at T₁ and the greatest girth (4.21 cm²) at T₄. There was no discernible change between the stem girth of T₁ and T₂ ($p > 0.05$). However, there was a significant difference between T₃ (2.79 cm²) and T₅ (4.21 cm²) ($p < 0.05$).

T₄ had the longest vines (76.03 cm), whereas T₁ had the shortest (57.87 cm). The length of the cucumber vine varied significantly across all treatments ($p < 0.05$). Even so, a statistical comparison of the cucumber vine length in T₄ and T₅ reveals no appreciable variation ($p > 0.05$). All of the treatments had

significantly varied leaf areas ($p < 0.05$), with T₄ producing the broadest (195 cm²) and T₁ producing the smallest (149.35 cm²) leaves. The number of leaves generated varies depending on the amount of cattle manure applied; T₄ reported the largest number of leaves (22.49) and T₁ the lowest (12.50). However, there was no noticeable difference generated ($p > 0.05$) between T₁ (12.50) and T₂ in terms of the number of leaves (Table 3).

According to the result in Table 4 yield attributes for cucumbers grown in the study area after various rates of cattle manure application, T₁ produced the fewest fruits (2.33), while T₄ produced the most (9.83). However, there was no statistically significant difference in the number of fruits taken from T₂ (5.17), T₃ (6.50), and T₅ (7.00) ($p > 0.05$). The fruit's circumference ranged from 14.27 cm in treatment T₁ to 17.19 cm in treatment T₄. The fruit circumference in T₁, T₂, and T₃, however, was not significantly influenced by the rates of animal manure. The length of cucumber fruits obviously varies with the amount of cattle manure applied, with T₄ producing the longest fruits (17.19 cm) and T₁ producing the smallest (7.68 cm). Although it was observed that treatments T₁ and T₂ had no effect on fruit length, T₄ recorded the lowest fruit weight (0.32 kg), and treatments T₁ and T₂ had the highest (5.14 kg). While T₄ and T₅ considerably influenced the fruit weight, the weight of the fruit recorded at T₁, T₂, and T₃ was found not to be significantly inclined ($p > 0.05$) to the application of cattle manure.

Discussion

The majority of the vital plant nutrients were insufficient in the experimental soil, suggesting that the soil lacks the fundamental elements needed for good crop growth and output. According to Adesina, et al. [14], the low soil fertility was caused by extensive and continuous cropping without any documented soil amendment methods to maintain its

Table 3: Effect of Varying Rates of Cattle Manure on the Vegetative Growth of Cucumber.

Treatments	Vine Girth (cm)	Vine Length (cm)	Leaf Area (cm ²)	Number of Leaves
T ₁ (0tha ⁻¹)	1.02 ± 0.64 ^a	57.87 ± 4.67 ^a	149.35 ± 2.98 ^a	12.50 ± 1.86 ^a
T ₂ (5tha ⁻¹)	1.45 ± 0.89 ^a	64.78 ± 3.64 ^{ab}	158.68 ± 5.17 ^b	13.79 ± 3.31 ^a
T ₃ (10tha ⁻¹)	2.79 ± 1.99 ^b	69.03 ± 4.54 ^b	156.68 ± 3.97 ^{ab}	16.24 ± 2.80 ^{ab}
T ₄ (15tha ⁻¹)	4.21 ± 1.42 ^c	76.03 ± 3.38 ^c	195.38 ± 2.48 ^d	22.49 ± 1.82 ^b
T ₅ (20tha ⁻¹)	3.45 ± 0.52 ^{cb}	73.46 ± 5.04 ^{bc}	171.22 ± 3.69 ^c	16.83 ± 3.20 ^{ab}

*Means with the same superscript in a column are not significantly different ($p > 0.05$).

Table 4: Effect of Varying Rates of Cattle Manure on the Yield of Cucumber.

Treatments	Number of Fruits	Fruit Circumference (cm)	Fruit Length (cm)	Fruit Weight (Kg)
T ₁ (0tha ⁻¹)	2.33 ± 0.97 ^a	14.27 ± 2.13 ^a	7.68 ± 1.12 ^a	0.32 ± 1.12 ^a
T ₂ (5tha ⁻¹)	5.17 ± 1.03 ^{ab}	15.62 ± 2.02 ^a	9.52 ± 2.09 ^a	0.80 ± 1.03 ^a
T ₃ (10tha ⁻¹)	6.50 ± 2.34 ^{ab}	15.49 ± 3.41 ^a	14.86 ± 4.42 ^c	1.22 ± 0.88 ^a
T ₄ (15tha ⁻¹)	9.83 ± 3.71 ^b	21.89 ± 5.98 ^c	17.19 ± 2.36 ^d	5.14 ± 2.33 ^c
T ₅ (20tha ⁻¹)	7.00 ± 1.93 ^{ab}	16.41 ± 3.55 ^{ab}	12.69 ± 1.18 ^{bc}	3.21 ± 1.47 ^b

*Means with the same superscript in a column are not significantly different ($p > 0.05$).

productivity. Soil fertility needs to be increased by the use of fertilizers in order to sustainably cultivate crops. The majority of soil physiochemical properties, such as soil fertility and lower soil acidity would be improved by the presence of essential plant nutrients in the cattle manure in a significant amount (Atmojo, 2003).

A lack of crucial nutrients may be the cause of the cucumber's poor performance in control plots when compared to those receiving cattle manure, which led to low yield and poor vegetative growth. The growth and output of cucumbers significantly increased when the rates of cattle dung were increased. The linear relationship between increases in cattle manure rates and vegetative growth and yield may be explained by the increased availability of nutrients that are more easily absorbed by the plants, resulting in increased production of photosynthates and efficient translocation for the development of the vegetative and reproductive parts, which in turn leads to higher fruit yields. This backs up the conclusions reached by Anjad, et al. [17] and Adesina, et al. [14], who believed that the usage of organic fertilizer improved the performance of maize and okra, respectively, in the treated plots compared to the untreated plots.

Due to the residual nutrient content of the soil being unsuitable for supporting cucumber plant growth, treatment 0 t ha⁻¹ (i.e., without cattle manure) could not support the proper growth of the plants. This resulted in the plants producing shorter vine lengths [18] and poor crop performance. However, plants that received 20 t ha⁻¹ of cattle manure provided sufficient nutrients to maintain the test plant's proper nutrition and growth. As a result, the crop has grown and developed more due to these plants' capacity for photosynthetic growth, resulting in the creation of longer vines per plant.

The maximum mean was found at T₄ (15 tha⁻¹), while the lowest mean was found at 0 t ha⁻¹ (i.e., without cattle manure). There was a substantial difference in vegetative growth between the various rates of cattle manure. The lowest mean recorded in the 0 t ha⁻¹ could be attributed to the fact that it could not support appropriate plant growth because the soil's residual nutrient content was inappropriate to support cucumber growth; likely, the nutrient content of the soil was below the critical level, which resulted in poor crop performance. In other words, the more nutrients added to the soil, the more leaves, vine length, vine girth, and leaf area there recorded. This outcome is in line with observations made by Ikeh, et al. [19], Hamma, et al. [18], Musara, and Chitamba [20], who found that cucumbers grown in soil that had been supplemented with manure had the highest growth parameters.

The role that cattle manure played in improving soil nutrients, and physical and chemical conditions, could be attributed to the significant improvement in growth attributes and yield parameters such as vine length, leaf area index, vine girth, number of leaves, fruit circumference, fruit length, number of fruits, and fruit weight. These improvements in turn led to an increase in growth rate and yield. As an organic component of the soil, cow dung helps to improve soil texture and bind water so that plant roots may more easily absorb

nutrients. The conclusions of this study were consistent with those of Fefiani and Barus [21], who found that cow manure significantly increased the length of the plant and the number of leaves on cucumbers. They were also in line with Ako's [22] claim that cow manure could increase the leaf area index in the growth of sorghum plants.

The availability of better nutrients, improved plant growth, and increased leaf production as a result of the positive effects of cattle manure on the physical and chemical properties of the soil may have made it possible for cucumber fruit yield attributes to significantly increase. Nevertheless, in spite of the obvious correlation between the rate of improvement in cucumber growth and the rate of application of cattle manure. The cucumber's performance started to deteriorate at T₅ (20 tha⁻¹), which contradicted earlier research from Enujeke [23] and Musara and Chitamba [20] that claimed cucumber plants that received 20 t ha⁻¹ of poultry manure experienced greater vegetative growth and fruit mass.

Due to the experimental site's poor soil fertility, the application of cattle dung had a considerable advantage over the control treatment. According to Ndaeyo, et al. [24,25], nutrient reserve in the soil influences crop response to fertilizer application. Crops respond more favorably to fertilizer application in soil with very low nutrient content than in soil with substantial nutrient reserve.

Conclusion

According to the current study, applying cattle manure greatly enhances cucumber growth and productivity. Therefore, it is crucial that farmers in the research region amend their soils with 15 tha⁻¹ cattle manure in order to get a higher fruit production. This would significantly reduce over-dependence on inorganic fertilizer for crop production and the amount of environmental pollution brought on by the disposal of livestock dung, which is typically thought of as a waste product in livestock enterprises would be significantly reduced.

References

1. Eifediyi EK, Remison SU. Growth and yield of cucumber (*Cucumis sativum* L) as influenced by farm yard manure and inorganic fertilizer. *Journal of Plant Breeding and Crop Science*. 2010; 2(7): 216-220.
2. Nweke IA, Orji EC, Ijearu SI. Effect of Staking and Plant Spacing on the Growth and Yield of Cucumber. *IOSR Journal of Environmental Science, Toxicology and Food Technology*. 2013; 3(4):26-31.
3. Mukherjee PK, Nema NK, Maity N, Sarkar BK. Phytochemical and therapeutic potential of cucumber. *Fitoterapia*. 2013 Jan;84:227-36. doi: 10.1016/j.fitote.2012.10.003. Epub 2012 Oct 23. PMID: 23098877.
4. Eifediyi EK, Remison SU. The effects of Inorganic Fertilizers on Yield of two varieties of Cucumber (*Cucumis sativus* L.) Report and Opinion. 2009; 1(5):74-80.
5. Wang X, Yan J, Zhang X, Zhang S, Chen Y. Organic manure input improves soil water and nutrients use for sustainable maize (*Zea mays* L) productivity on the Loess Plateau. *PLoS One*. 2020 Aug 25;15(8):e0238042. doi: 10.1371/journal.pone.0238042. PMID: 32841280; PMCID: PMC7447034.
6. Deksisssa T, Short I, Allen J. Effect of soil amendment with compost on growth and water use efficiency of Amaranth. In *Int. Water Resources: Challenges for*



- the 21st Century & Water Resources Education, Proc. UCOWR/NIWR Annual Conf. July 22-24, 2008; Durham NC.
7. John LW, Jamer DB, Samuel LT, Waner LW. Soil Fertility and Fertilizer: An Introduction to Nutrient Management. Pearson Education, India. 2004; 106-153.
 8. Maclean EO Aluminum: In Black, CA. (eds). Methods of Soil Analysis, Part 2. Agro., American Soc. of Agro., Madison, Wisconsin, USA. 1982.
 9. Gee GW, Bauder JW. Particle size analysis. Methods of soil Analysis, Part 1: In Klute, A. (eds) physical and mineralogical methods. Agronomy Monograph (9). Agro. and Soil Sc. Society of America. Madison, Wisconsin. 1986.
 10. Bremner JM, Mulvaney CS. Total nitrogen determination. 595-622. In: methods of soil analysis part 2 chemical microbiological properties. 2nd edition. Agronomy monograph. 1982; 9.
 11. Thomas GW. Exchangeable cations. In: Page A. L, et al., (eds). Methods of Soil Analysis part 2: 2nd edition Agronomy monograph no.9, Chemical and microbiological properties. American Soc. of Agro. Madison, Wisconsin. 1982; 159-165.
 12. International Institute for Tropical Agriculture (IITA). Laboratory manual for soil and plant analysis. Manual series 7. IITA. Ibadan. Nigeria. 1979.
 13. Olsen SR, Sommers LE. Determination of available phosphorus, pp 403-430. In: Methods of soil analysis part 2. Chemical and Micro. Properties. Agro. Mono. No. 9 (2nd edition). 1994.
 14. Adesina JM, Ileke KD, Adebayo RA. *Podagrira unifroma* and *Nisota dilecta* (Coleoptera: Chrysomalidae) Infestation on Okra (*Abelmoschus esculentus*) in Response to Nutrient Sources. Journal of Crop Nutrition Science. 2018; 4(2): 45-55.
 15. Tahir SM, Usman IS, Katung MD, Ishiyaku MF. In Situ Germination and Early Seedling Growth of Wormwood (*Artemisia annua* L.) American Journal of Plant Sciences. 2014; 5:1694-1701.
 16. Statistical Analysis System. SAS User's Guide Statistic Version 5 SAS Institute Inc. Raleigh North California, U.S.A. 2010; 956.
 17. Anjad A, Ali F, Hue NV. Nitrogen Synchronization from organic manure, Application as measured from the soil solution, and SPAD readings for sweet corn. 2003; 2.
 18. Hamma IL, Ibrahim U, Haruna M. Effect of poultry manure on the growth and yield of cucumber (*Cucumis sativum* L.) in Samaru, Zaria. Nigeria Journal of Agriculture, Food, and Environment. 2012; 8(1):94-98.
 19. Ikeh AO, Udoh EI, Uduak GI, Udounang PI, Etokeren UE. Response of cucumber (*Cucumis sativus* L.) to different rates of goat and poultry manure on an ultisol. Journal of Agriculture & Social Research. 2012; 12(2):132-139.
 20. Musara C, Chitamba J. Evaluation of cattle manure application rate on the growth rate and fruit yield of cucumber (*Cucumis sativa* L.). International Research Journal of Agric Science & Soil Science. 2014; 4(9):167-171.
 21. Fefiani Y, Barus WA. Respon Pertumbuhan dan Produksi Tanaman Mentimun (*Cucumis sativus* L.) Akibat Pemberian Cow manure Dan Pupuk Organik Padat Supernasa. Agrium. 2014; 19(1):21-30.
 22. Ako A. Pengaruh Tingkat Pemberian Pupuk Kandang terhadap Pertumbuhan dan Produksi Rumpun Gajah. Media Veteriner. 1997;4 (2):34-43.
 23. Enujoke E. Growth and yield responses of cucumber to five different rates of poultry manure in the Asaba area of Delta State, Nigeria. International Research Journal of Agricultural Science and Soil Science. 2013; 3(11):369-375.
 24. Ndaeyo NU, Ukpong ES, John NM. Performances of Okra as affected by organic and inorganic fertilizers on util. Proceedings of the 39th Conference of the Agricultural Society of Nigeria October 9-13, 2005; 206-209.
 25. Walkley A, Black IA. An examination of the Degtjareff method for determining soil organic matter and proposed modification of the Chromic acid titration method. Soil Sc. 1934; 37:29-38.

Discover a bigger Impact and Visibility of your article publication with Peertechz Publications

Highlights

- ❖ Signatory publisher of ORCID
- ❖ Signatory Publisher of DORA (San Francisco Declaration on Research Assessment)
- ❖ Articles archived in worlds' renowned service providers such as Portico, CNKI, AGRIS, TDNet, Base (Bielefeld University Library), CrossRef, Scilit, J-Gate etc.
- ❖ Journals indexed in ICMJE, SHERPA/ROME0, Google Scholar etc.
- ❖ OAI-PMH (Open Archives Initiative Protocol for Metadata Harvesting)
- ❖ Dedicated Editorial Board for every journal
- ❖ Accurate and rapid peer-review process
- ❖ Increased citations of published articles through promotions
- ❖ Reduced timeline for article publication

Submit your articles and experience a new surge in publication services (<https://www.peertechz.com/submission>).

Peertechz journals wishes everlasting success in your every endeavours.